arXiv:1410.4153v3 [math.NT] 27 Sep 2016

On quintic identities

By Kazuhide Matsuda

Faculty of Fundamental Science, National Institute of Technology, Niihama College,
7-1 Yagumo-chou, Niihama, Ehime, 792-8580, Japan
e-mail: matsuda@sci.niihama-nct.ac.jp

Abstract In this paper, we derive quintic versions of the cubic identities of Farkas and Kra. In particular, we believe that our results can be easily generalized to k th power versions, $(k = 7, 9, 11, \ldots)$. Moreover, we investigate the algebraic structure of theta constants of level five.
Key words theta functions; theta constants; rational characteristic; resultant
MSC(2010) 14K25

1 Introduction

Throughout this paper, let $\mathbb{N}$, $\mathbb{N}_0$, $\mathbb{Z}$, $\mathbb{Q}$, $\mathbb{R}$ and $\mathbb{C}$ denote the set of positive integers, nonnegative integers, integers, quotient numbers, real numbers and complex numbers, respectively. Moreover, let the upper half plane be defined by

$$\mathbb{H}^2 = \{\tau \in \mathbb{C} \,|\, \Im\tau > 0\}.$$

Following Farkas and Kra [2], we introduce the theta function with characteristic $\begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} \in \mathbb{R}^2$, which is defined by

$$\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau) = \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta) := \sum_{n \in \mathbb{Z}} \exp\left(2\pi i \left[\frac{1}{2}\left(n + \frac{\epsilon}{2}\right)^2 \tau + \left(n + \frac{\epsilon}{2}\right)\left(\zeta + \frac{\epsilon'}{2}\right)\right]\right),$$

which uniformly and absolutely converges on compact subsets of $\mathbb{C} \times \mathbb{H}^2$, where $\mathbb{H}^2$ is the upper half-plane. The theta constants are defined by

$$\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} := \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (0, \tau).$$

Farkas and Kra [2] treated the theta constants with rational characteristics, that is, the case where ϵ and ϵ' are both rational numbers, and derived a number of interesting theta constant identities.

One of the most famous theta constant identities is Jacobi's quartic, which is given by

$$\theta^4\left[\begin{array}{c}0\\0\end{array}\right]=\theta^4\left[\begin{array}{c}1\\0\end{array}\right]+\theta^4\left[\begin{array}{c}0\\1\end{array}\right].$$

Based on the uniformization theory, Farkas and Kra [1] proved the following cubic identities: for every $\tau\in\mathbb{H}^2$,

$$\theta^3\left[\begin{array}{c}\frac{1}{3}\\\frac{1}{3}\end{array}\right]+\theta^3\left[\begin{array}{c}\frac{1}{3}\\\frac{5}{3}\end{array}\right]=\theta^3\left[\begin{array}{c}\frac{1}{3}\\1\end{array}\right], \tag{1.1}$$

and

$$\exp\left(\frac{\pi i}{3}\right)\theta^3\left[\begin{array}{c}\frac{1}{3}\\\frac{1}{3}\end{array}\right]+\exp\left(\frac{2\pi i}{3}\right)\theta^3\left[\begin{array}{c}\frac{1}{3}\\\frac{5}{3}\end{array}\right]=\theta^3\left[\begin{array}{c}1\\\frac{1}{3}\end{array}\right]. \tag{1.2}$$

Our aim is to obtain quintic versions of the identities (1.1) and (1.2). Moreover, we investigate the algebraic structure of the following theta constants:

$$\theta\left[\begin{array}{c}\frac{j}{5}\\\frac{k}{5}\end{array}\right],\ \theta\left[\begin{array}{c}1\\\frac{j}{5}\end{array}\right],\ (j=1,3,k=1,3,5,7,9).$$

This paper is organized as follows. In Section 2, we review the properties of the theta functions. In Section 3, we derive quintic versions of the identities (1.1) and (1.2). In particular, we believe that our results can be easily generalized to k th power versions, $(k=7,9,11,\ldots)$.

In Section 4, we prove that $\theta\left[\begin{array}{c}1\\1/5\end{array}\right]/\theta\left[\begin{array}{c}1\\3/5\end{array}\right]$ is given by a rational expression of the following theta constants:

$$\theta\left[\begin{array}{c}\frac{j}{5}\\\frac{k}{5}\end{array}\right],(j=1,3,k=1,3,5,7,9). \tag{1.3}$$

In Section 5, we show that $\theta\left[\begin{array}{c}1\\1/5\end{array}\right]$, or $\theta\left[\begin{array}{c}1\\3/5\end{array}\right]$ is given by rational expessions of the theta constants (1.3).

In Section 6, for $j=1,3$, we prove that one of $\theta\left[\begin{array}{c}j/5\\k/5\end{array}\right]$, $(k=1,3,5,7,9)$, is given by two rational expressions of four of the following theta constants:

$$\theta\left[\begin{array}{c}\frac{j}{5}\\\frac{k}{5}\end{array}\right],(k=1,3,5,7,9).$$

For this purpose, considering the determinant structure, we obtain the theta constant identities. This method is based on the method of Matsuda [4].

In Section 7, for $j=1,3$, we derive the theta constant identities among $\theta\left[\begin{array}{c}j/5\\k/5\end{array}\right]$, $(k=1,3,5,7,9)$. In Section 8, we investigate the algebraic structure of the theta constants (1.3).

Remark

By the identity (1.1), Farkas [3] showed that for each nonnegative integer $n \in \mathbb{N}_0$,

$$\sigma(3n+2) = 3\sum_{k=0}^{n} \delta(3k+1)\delta(3(n-k)+1),$$

where for each positive integer $n \in \mathbb{N}$,

$$\sigma(n) = \sum_{d|n} d, \text{ and } \delta(n) = d_{1,3}(n) - d_{2,3}(n).$$

We believe that the theta constant identities of this paper can be applied to number theory. However, the computation is more complicated.

Acknowledgments

We are grateful to Professor H. Watanabe for his useful comments.

2 The properties of the theta functions

We first note that for $m, n \in \mathbb{Z}$,

$$\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta + n + m\tau, \tau) = \exp(2\pi i) \left[\frac{n\epsilon - m\epsilon'}{2} - mz - \frac{m^2\tau}{2} \right] \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau), \tag{2.1}$$

and

$$\theta \begin{bmatrix} \epsilon + 2m \\ \epsilon' + 2n \end{bmatrix} (\zeta, \tau) = \exp(\pi i \epsilon n) \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau). \tag{2.2}$$

Furthermore,

$$\theta \begin{bmatrix} -\epsilon \\ -\epsilon' \end{bmatrix} (\zeta, \tau) = \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (-\zeta, \tau) \text{ and } \theta' \begin{bmatrix} -\epsilon \\ -\epsilon' \end{bmatrix} (\zeta, \tau) = -\theta' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (-\zeta, \tau).$$

For $m, n \in \mathbb{R}$, we see that

$$\begin{aligned} &\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} \left(\zeta + \frac{n + m\tau}{2}, \tau \right) \\ &= \exp(2\pi i) \left[-\frac{m\zeta}{2} - \frac{m^2\tau}{8} - \frac{m(\epsilon' + n)}{4} \right] \theta \begin{bmatrix} \epsilon + m \\ \epsilon' + n \end{bmatrix} (\zeta, \tau). \end{aligned} \tag{2.3}$$

We note that $\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](\zeta,\tau)$ has only one zero in the fundamental parallelogram, which is given by

$$\zeta=\frac{1-\epsilon}{2}\tau+\frac{1-\epsilon'}{2}.$$

All the theta functions have infinite product expansions, which are given by

$$\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](\zeta,\tau)=\exp\left(\frac{\pi i\epsilon\epsilon'}{2}\right)x^{\frac{\epsilon^2}{4}}z^{\frac{\epsilon}{2}}$$
$$\prod_{n=1}^{\infty}(1-x^{2n})(1+e^{\pi i\epsilon'}x^{2n-1+\epsilon}z)(1+e^{-\pi i\epsilon'}x^{2n-1-\epsilon}/z), \tag{2.4}$$

where $x=\exp(\pi i\tau)$, $z=\exp(2\pi i\zeta)$.

Following Farkas and Kra [2], we last define $\mathcal{F}_N\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right]$ to be the set of the entire functions f that satisfy the two functional equations,

$$f(\zeta+1)=\exp(\pi i\epsilon)\ f(\zeta),$$

and

$$f(\zeta+\tau)=\exp(-\pi i)[\epsilon'+2N\zeta+N\tau]\ f(\zeta),\quad \zeta\in\mathbb{C},\ \tau\in\mathbb{H}^2,$$

where N is a positive integer and $\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right]\in\mathbb{R}^2$. This set of functions is called the space of *N-th order θ-functions with characteristic* $\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right]$. Note that

$$\dim\mathcal{F}_N\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right]=N.$$

For its proof, see Farkas and Kra [2, pp.131].

3 Quintic identities

Theorem 3.1. *For every $\tau\in\mathbb{H}^2$, we have*

$$\theta^5\left[\begin{array}{c}\frac{1}{5}\\ \frac{1}{5}\end{array}\right]-\theta^5\left[\begin{array}{c}\frac{1}{5}\\ \frac{3}{5}\end{array}\right]+\theta^5\left[\begin{array}{c}\frac{1}{5}\\ 1\end{array}\right]-\theta^5\left[\begin{array}{c}\frac{1}{5}\\ \frac{7}{5}\end{array}\right]+\theta^5\left[\begin{array}{c}\frac{1}{5}\\ \frac{9}{5}\end{array}\right]=0, \tag{3.1}$$

and

$$\theta^5\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]-\theta^5\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]+\theta^5\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]-\theta^5\left[\begin{array}{c}\frac{3}{5}\\ \frac{7}{5}\end{array}\right]+\theta^5\left[\begin{array}{c}\frac{3}{5}\\ \frac{9}{5}\end{array}\right]=0. \tag{3.2}$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^5 \begin{bmatrix} 1 \\ 1 \end{bmatrix}(z)}{\theta \begin{bmatrix} \frac{1}{5} \\ \frac{1}{5} \end{bmatrix}(z)\theta \begin{bmatrix} \frac{1}{5} \\ \frac{3}{5} \end{bmatrix}(z)\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}(z)\theta \begin{bmatrix} \frac{1}{5} \\ \frac{7}{5} \end{bmatrix}(z)\theta \begin{bmatrix} \frac{1}{5} \\ \frac{9}{5} \end{bmatrix}(z)},$$

and

$$\psi(z) = \frac{\theta^5 \begin{bmatrix} 1 \\ 1 \end{bmatrix}(z)}{\theta \begin{bmatrix} \frac{3}{5} \\ \frac{1}{5} \end{bmatrix}(z)\theta \begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}(z)\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}(z)\theta \begin{bmatrix} \frac{3}{5} \\ \frac{7}{5} \end{bmatrix}(z)\theta \begin{bmatrix} \frac{3}{5} \\ \frac{9}{5} \end{bmatrix}(z)}.$$

By $\varphi(z)$, we obtain equation (3.1). Equation (3.2) can be derived by $\psi(z)$ in the same way.

We first note that in the fundamental parallelogram, the poles of $\varphi(z)$ are $z = (2\tau + 2)/5, (2\tau + 1)/5, 2\tau/5, (2\tau - 1)/5$, and $(2\tau - 2)/5$. The direct calculation yields

$$\operatorname{Res}\left(\varphi(z), \frac{2\tau+2}{5}\right) = -\frac{\theta^5 \begin{bmatrix} \frac{1}{5} \\ \frac{1}{5} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}},$$

$$\operatorname{Res}\left(\varphi(z), \frac{2\tau+1}{5}\right) = \frac{\theta^5 \begin{bmatrix} \frac{1}{5} \\ \frac{3}{5} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}},$$

$$\operatorname{Res}\left(\varphi(z), \frac{2\tau}{5}\right) = -\frac{\theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}},$$

$$\operatorname{Res}\left(\varphi(z), \frac{2\tau-1}{5}\right) = \frac{\theta^5 \begin{bmatrix} \frac{1}{5} \\ \frac{7}{5} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}},$$

and

$$\operatorname{Res}\left(\varphi(z), \frac{2\tau-2}{5}\right) = -\frac{\theta^5 \begin{bmatrix} \frac{1}{5} \\ \frac{9}{5} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}\theta^2 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}.$$

Since

$$\begin{aligned}\operatorname{Res}(\varphi(z),(2\tau+2)/5)+\operatorname{Res}(\varphi(z),(2\tau+1)/5)+\operatorname{Res}(\varphi(z),2\tau/5)\\ +\operatorname{Res}(\varphi(z),(2\tau-1)/5)+\operatorname{Res}(\varphi(z),(2\tau-2)/5)=0,\end{aligned}$$

equation (3.1) follows. □

Theorem 3.2. *For every $\tau\in\mathbb{H}^2$, we have*

$$\zeta_5\theta^5\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}+\zeta_5^3\theta^5\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}+\theta^5\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}-\zeta_5^2\theta^5\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}-\zeta_5^4\theta^5\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}=0, \tag{3.3}$$

and

$$\zeta_5^3\theta^5\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}+\zeta_5^4\theta^5\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}+\theta^5\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}-\zeta_5\theta^5\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}-\zeta_5^2\theta^5\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}=0. \tag{3.4}$$

Proof. Consider the following elliptic functions:

$$\varphi(z)=\frac{\theta^5\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{7}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{9}{5}\\ \frac{1}{5}\end{bmatrix}(z)},$$

and

$$\psi(z)=\frac{\theta^5\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{7}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{9}{5}\\ \frac{3}{5}\end{bmatrix}(z)}.$$

The theorem can be proved in the same way as Theorem 3.1. □

Rermark

The identity (3.1) was proved by Farkas and Kra [2, pp. 277]. We believe that the other identities in Theorems 3.1 and 3.2 are new.

4 Rational expressions of theta constants (1)

4.1 The case for $j=1$

In this subsection, we prove the following theorem:

Theorem 4.1. $\theta\begin{bmatrix}1\\1/5\end{bmatrix}/\theta\begin{bmatrix}1\\3/5\end{bmatrix}$ *is given by a rational expression of four of* $\theta\begin{bmatrix}1/5\\k/5\end{bmatrix}$, *where* $k=1,3,5,7,9$.

For the proof, we obtain the following proposition and theorem:

Proposition 4.2. *For every* $(\zeta,\tau)\in\mathbb{C}\times\mathbb{H}^2$, *we have*

$$\begin{aligned}&\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)+\zeta_5^2\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{9}{5}\end{bmatrix}(\zeta,\tau)\\&-\zeta_5^4\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau)-\zeta_5^2\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)=0,\end{aligned}\tag{4.1}$$

$$\begin{aligned}&\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\1\end{bmatrix}(\zeta,\tau)-\zeta_5^2\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\1\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau)\\&+\zeta_5^2\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)-\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)=0,\end{aligned}\tag{4.2}$$

$$\begin{aligned}&\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau)-\zeta_5^2\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\1\end{bmatrix}(\zeta,\tau)\\&+\zeta_5^2\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\1\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)+\zeta_5^2\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)=0,\end{aligned}\tag{4.3}$$

$$\begin{aligned}&\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)+\zeta_5^2\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\\&-\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\1\end{bmatrix}(\zeta,\tau)+\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\1\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)=0,\end{aligned}\tag{4.4}$$

$$\begin{aligned}&\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau)-\zeta_5^2\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\\&+\zeta_5^2\theta\begin{bmatrix}1\\\frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\1\end{bmatrix}(\zeta,\tau)-\theta\begin{bmatrix}1\\\frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\1\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)=0.\end{aligned}\tag{4.5}$$

Proof. We treat equation (4.1). The others can be proved in the same way. For this purpose, we first note that $\dim\mathcal{F}_3\begin{bmatrix}\frac{3}{5}\\1\end{bmatrix}=3$ and

$$\begin{aligned}&\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau),\ \theta^2\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{3}{5}\end{bmatrix}(\zeta,\tau)\\&\theta^2\begin{bmatrix}\frac{1}{5}\\\frac{9}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau),\ \theta^2\begin{bmatrix}\frac{1}{5}\\\frac{7}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\\frac{1}{5}\end{bmatrix}(\zeta,\tau)\in\mathcal{F}_3\begin{bmatrix}\frac{3}{5}\\1\end{bmatrix}.\end{aligned}$$

Therefore, there exist complex numbers, x_1, x_2, x_3, x_4, not all zero such that

$$x_1\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)+x_2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)$$
$$+x_3\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)+x_4\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)=0.$$

By substituting

$$\zeta=\frac{2\tau\pm 2}{5},\frac{2\tau\pm 1}{5},$$

we have the following system of equations:

$$\begin{aligned}
x_2\zeta_5^2\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}+x_3\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix} &= 0,\\
x_3\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}-x_4\zeta_5^2\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix} &= 0,\\
x_1\zeta_5^2\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}+x_4\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix} &= 0,\\
x_1\zeta_5^2\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}-x_2\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix} &= 0.
\end{aligned}$$

Solving this system of equations, we obtain

$$(x_1,x_2,x_3,x_4)=\alpha\left(\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix},\zeta_5^2\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix},-\zeta_5^4\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix},-\zeta_5^2\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\right),$$

for some nonzero complex number $\alpha\in\mathbb{C}^*$, which proves the proposition. □

Theorem 4.3. *For every $\tau\in\mathbb{H}^2$, we have*

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)-\zeta_5^4\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)}{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)-\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)},\tag{4.6}$$

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)+\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)+\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)},\tag{4.7}$$

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)-\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}(0,\tau)+\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)},\tag{4.8}$$

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{-\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)+\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)}{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)+\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)},\tag{4.9}$$

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)+\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)}{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}(0,\tau)+\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)}.\tag{4.10}$$

Proof. We deal with equation (4.6). The others can be proved in the same way. For this purpose, setting $\zeta=0$ in equation (4.1), we have

$$\begin{aligned}&\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)\left\{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)-\zeta_5^4\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}(0,\tau)\right\}\\ =&\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)\left\{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)-\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\right\}.\end{aligned}$$

Suppose that for every $\tau\in\mathbb{H}^2$,

$$\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)-\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\equiv 0,$$

which implies that

$$\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)-\zeta_5^4\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}(0,\tau)\equiv 0.$$

Thus, we have

$$\zeta_5^2-\left(\frac{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)}\right)^5\equiv 0,$$

which implies that $\theta\left[\begin{array}{c}1/5\\9/5\end{array}\right](0,\tau)/\theta\left[\begin{array}{c}1/5\\1/5\end{array}\right](0,\tau)$ is a constant. Taking the limit $\tau\longrightarrow i\infty$, we obtain

$$\theta\left[\begin{array}{c}\frac{1}{5}\\\frac{9}{5}\end{array}\right](0,100\tau)=\exp\left(\frac{8\pi i}{50}\right)\theta\left[\begin{array}{c}\frac{1}{5}\\\frac{1}{5}\end{array}\right](0,100\tau),$$

where we used the Jacobi triple product identity (2.4). Since

$$\theta\left[\begin{array}{c}\frac{1}{5}\\\frac{9}{5}\end{array}\right](0,100\tau)=\sum_{n\in\mathbb{Z}}\exp(2\pi i)\left[\left(n+\frac{1}{10}\right)\frac{9}{10}\right]x^{(10n+1)^2},$$

$$\theta\left[\begin{array}{c}\frac{1}{5}\\\frac{1}{5}\end{array}\right](0,100\tau)=\sum_{n\in\mathbb{Z}}\exp(2\pi i)\left[\left(n+\frac{1}{10}\right)\frac{1}{10}\right]x^{(10n+1)^2},\ x=\exp(\pi i\tau),$$

comparing the coefficients of x^{121}, we obtain $\zeta_5^3=1$, which is impossible.

Thus, it follows that

$$\zeta_5^2\theta^2\left[\begin{array}{c}\frac{1}{5}\\\frac{7}{5}\end{array}\right](0,\tau)\theta\left[\begin{array}{c}\frac{1}{5}\\\frac{1}{5}\end{array}\right](0,\tau)-\zeta_5^2\theta^2\left[\begin{array}{c}\frac{1}{5}\\\frac{3}{5}\end{array}\right](0,\tau)\theta\left[\begin{array}{c}\frac{1}{5}\\\frac{3}{5}\end{array}\right](0,\tau)\not\equiv 0,$$

which proves the theorem. □

4.2 The case for $j=3$

In this subsection, we prove the following theorem:

Theorem 4.4. *$\theta\left[\begin{array}{c}1\\1/5\end{array}\right]/\theta\left[\begin{array}{c}1\\3/5\end{array}\right]$ is given by a rational expression of four of $\theta\left[\begin{array}{c}3/5\\k/5\end{array}\right]$, where $k=1,3,5,7,9$.*

For the proof, we show the following proposition and theorem, which can be proved in the same way as in the previous subsection.

Proposition 4.5. *For every $(\zeta,\tau)\in\mathbb{C}\times\mathbb{H}^2$, we have*

$$\begin{aligned}&\theta\left[\begin{array}{c}1\\\frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right](\zeta,\tau)+\zeta_5\theta\left[\begin{array}{c}1\\\frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{9}{5}\end{array}\right](\zeta,\tau)\\&-\zeta_5^2\theta\left[\begin{array}{c}1\\\frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{9}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{7}{5}\end{array}\right](\zeta,\tau)-\zeta_5\theta\left[\begin{array}{c}1\\\frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{7}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right](\zeta,\tau)=0,\end{aligned}\tag{4.11}$$

$$\begin{aligned}&\theta\left[\begin{array}{c}1\\\frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right](\zeta,\tau)-\zeta_5\theta\left[\begin{array}{c}1\\\frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{7}{5}\end{array}\right](\zeta,\tau)\\&+\zeta_5\theta\left[\begin{array}{c}1\\\frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{7}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right](\zeta,\tau)-\theta\left[\begin{array}{c}1\\\frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right](\zeta,\tau)=0,\end{aligned}\tag{4.12}$$

$$\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)-\zeta_5\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(\zeta,\tau)$$
$$+\zeta_5\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)+\zeta_5\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)=0, \tag{4.13}$$

$$\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)+\zeta_5\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)$$
$$-\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(\zeta,\tau)+\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)=0, \tag{4.14}$$

$$\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)-\zeta_5\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)$$
$$+\zeta_5\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(\zeta,\tau)-\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)=0. \tag{4.15}$$

Theorem 4.6. *For every $\tau\in\mathbb{H}^2$, we have*

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)-\zeta_5^2\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)}{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)-\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)}, \tag{4.16}$$

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)+\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(0,\tau)+\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)}, \tag{4.17}$$

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(0,\tau)-\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)+\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)}, \tag{4.18}$$

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{-\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)+\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(0,\tau)}{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)+\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(0,\tau)}, \tag{4.19}$$

$$\frac{\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}(0,\tau)}=\frac{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)+\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)}{\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)+\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}(0,\tau)}. \tag{4.20}$$

5 Rational expressions of theta constants (2)

Theorem 5.1. *For every $\tau \in \mathbb{H}^2$, we have*

$$\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix} = \zeta_5^2 \frac{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix} + \zeta_5^2\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}}{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}}, \ \theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix} = \zeta_5 \frac{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix} - \zeta_5^2\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}}{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}}.$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(z)}, \ \psi(z) = \frac{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(z)\theta\begin{bmatrix}-\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(z)}.$$

By $\varphi(z)$, we derive the rational expression of $\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}$. The other can be obtained by $\psi(z)$ in the same way.

We first note that in the fundamental parallelogram, the poles of $\varphi(z)$ are $z = (2\tau - 2)/5$, $(2\tau+1)/5$ and $(\tau+1)/5$. The direct calculation yields

$$\mathrm{Res}\left(\varphi(z), \frac{2\tau-2}{5}\right) = -\zeta_5 \frac{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}}{\theta'\begin{bmatrix}1\\ 1\end{bmatrix}\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}},$$

$$\mathrm{Res}\left(\varphi(z), \frac{2\tau+1}{5}\right) = -\frac{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}}{\theta'\begin{bmatrix}1\\ 1\end{bmatrix}\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}},$$

and

$$\mathrm{Res}\left(\varphi(z), \frac{\tau+1}{5}\right) = \zeta_5^4 \frac{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}}{\theta'\begin{bmatrix}1\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}}.$$

Since $\mathrm{Res}\left(\varphi(z), (2\tau-2)/5\right) + \mathrm{Res}\left(\varphi(z), (2\tau+1)/5\right) + \mathrm{Res}\left(\varphi(z), (\tau+1)/5\right) = 0$, we can obtain the rational expression. □

Theorem 5.2. *For every $\tau \in \mathbb{H}^2$, we have*

$$\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix} = \zeta_5 \frac{\zeta_5\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix} + \theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}}{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}},\ \theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix} = \frac{-\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix} + \theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}}{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}}.$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(z)},\ \psi(z) = \frac{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}-\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(z)}.$$

The theorem can be proved in the same way as Theorem 5.1. □

Theorem 5.3. *For every $\tau \in \mathbb{H}^2$, we have*

$$\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix} = \zeta_5 \frac{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix} - \theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}}{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}},\ \theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix} = \zeta_5^4 \frac{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix} - \theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}}{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}}.$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{1}{5}\\ -\frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}(z)},\ \psi(z) = \frac{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(z)\theta\begin{bmatrix}-\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(z)}.$$

The theorem can be proved in the same way as Theorem 5.1. □

Theorem 5.4. *For every $\tau \in \mathbb{H}^2$, we have*

$$\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix} = -\zeta_5^2 \frac{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix} + \zeta_5^3\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}}{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}}, \theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix} = -\zeta_5^3 \frac{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix} - \theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}}{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{9}{5}\end{bmatrix}}.$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{1}{5}\\ -\frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}(z)},\ \psi(z) = \frac{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}(z)\theta\begin{bmatrix}-\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(z)}.$$

The theorem can be proved in the same way as Theorem 5.1. □

6 Rational expressions of theta constants (3)

6.1 Review of the resultant

Set

$$f(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n,$$
$$g(x) = b_0x^n + b_1x^{m-1} + \cdots + b_m.$$

We then define the resultant $R(f,g)$ by the following $(m+n)\times(m+n)$ determinant:

$$R(f,g) = \begin{vmatrix} a_0 & a_1 & \cdots & a_n & & & \mathbf{0} \\ & a_0 & a_1 & \cdots & a_n & & \\ \mathbf{0} & & \ddots & \ddots & & \ddots & \\ & & & a_0 & a_1 & \cdots & a_n \\ b_0 & b_1 & \cdots & b_m & & & \mathbf{0} \\ & b_0 & b_1 & \cdots & b_m & & \\ \mathbf{0} & & \ddots & \ddots & & \ddots & \\ & & & b_0 & b_1 & \cdots & b_m \end{vmatrix}.$$

$R(f,g)$ shows us the condition for $f(x)$ and $g(x)$ to have a common zero.

Proposition 6.1. *$f(x)$ and $g(x)$ both have a common zero if and only if $R(f,g)=0$.*

Especially, when

$$f(x) = a_0x^2 + a_1x + a_2, \text{ and } g(x) = b_0x^2 + b_1x + b_2,$$

the resultant is given by

$$\begin{aligned} R(f,g) &= \begin{vmatrix} a_0 & a_1 & a_2 & 0 \\ 0 & a_0 & a_1 & a_2 \\ b_0 & b_1 & b_2 & 0 \\ 0 & b_0 & b_1 & b_2 \end{vmatrix} \\ &= (a_0b_2 - a_2b_0)^2 - (a_0b_1 - a_1b_0)(a_1b_2 - a_2b_1). \end{aligned}$$

6.2 The case for $j = 1$

In this subsection, we prove the following theorem:

Theorem 6.2. *One of $\theta\left[\begin{smallmatrix} 1/5 \\ k/5 \end{smallmatrix}\right], (k = 1,3,5,7,9)$, is given by two rational expressions of the other four theta constants.*

For the proof, we obtain the following proposition and theorem:

Proposition 6.3. *For every* $(\zeta,\tau)\in\mathbb{C}\times\mathbb{H}^2$, *we have*

$$\theta^2\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)-\theta^2\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)$$
$$+\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(\zeta,\tau)=0, \tag{6.1}$$

$$\zeta_5^2\theta^2\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)-\zeta_5^2\theta^2\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)$$
$$+\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)=0, \tag{6.2}$$

$$\theta^2\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)+\zeta_5^2\theta^2\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)$$
$$-\zeta_5^2\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)=0, \tag{6.3}$$

$$\theta^2\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(\zeta,\tau)+\zeta_5^2\theta^2\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)$$
$$-\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)=0, \tag{6.4}$$

$$\theta^2\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(\zeta,\tau)-\theta^2\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(\zeta,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(\zeta,\tau)$$
$$+\zeta_5^2\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(\zeta,\tau)=0. \tag{6.5}$$

Proof. Equation (6.1) was proved by Matsuda [4], who proved it in the same way as Proposition 4.2. The others can be also proved in the same way. □

Theorem 6.4. *For every* $\tau\in\mathbb{H}^2$, *we have*

$$\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}=\frac{\left\{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}-\theta^3\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}\left\{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}+\zeta_5^3\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\right\}}{\left\{\zeta_5^3\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}+\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\right\}^2}, \tag{6.6}$$

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]=\frac{\left\{\zeta_5^2\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]+\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\right\}\left\{\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]-\zeta_5^4\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}}{\left\{\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]+\zeta_5^2\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}^2},\tag{6.7}$$

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]=\frac{\left\{\zeta_5^3\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]+\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}\left\{\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]+\zeta_5^3\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}}{\left\{\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]-\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\right\}^2},\tag{6.8}$$

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]=\frac{\left\{\zeta_5^2\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]+\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\right\}\left\{\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]-\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}}{\left\{\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]+\zeta_5^3\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\right\}^2},\tag{6.9}$$

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]=\frac{\left\{\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]-\zeta_5\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\right\}\left\{-\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]+\zeta_5^4\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}}{\left\{-\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]+\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\right\}^2},\tag{6.10}$$

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]=\frac{\left\{-\zeta_5^2\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]+\zeta_5^2\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\right\}\left\{-\zeta_5^4\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]+\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}}{\left\{\zeta_5^4\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]-\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}^2},\tag{6.11}$$

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]=\frac{\left\{\zeta_5^2\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]+\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\right\}\left\{\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]+\zeta_5^2\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]\right\}}{\left\{\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]-\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}^2},\tag{6.12}$$

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]=\frac{\left\{\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]+\zeta_5^3\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\right\}\left\{-\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]+\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\right\}}{\left\{\zeta_5^2\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]+\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\right\}^2},\tag{6.13}$$

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]=\frac{\left\{\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]-\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\right\}\left\{\zeta_5^2\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]+\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\right\}}{\left\{\zeta_5^2\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]+\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\right\}^2},\tag{6.14}$$

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]=\frac{\left\{\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]+\zeta_5^3\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\right\}\left\{-\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]+\zeta_5^4\theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\right\}}{\left\{\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{2}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]+\zeta_5^2\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{4}{5}\end{smallmatrix}\right]\right\}^2}.\tag{6.15}$$

Proof. By equations (6.1) and (6.3), we prove equation (6.6). The others can be proved by two equations of Proposition 6.3 in the same way.

Equations (6.1) and (6.3) show that for arbitrary complex numbers $z, w \in \mathbb{C}$,

$$\theta^2\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(z,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(z,\tau)-\theta^2\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(z,\tau)$$
$$+\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(z,\tau)=0,$$

and

$$\theta^2\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(w,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(w,\tau)+\zeta_5^2\theta^2\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(w,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(w,\tau)$$
$$-\zeta_5^2\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(w,\tau)=0,$$

which implies that $\theta\begin{bmatrix}1\\ \frac{1}{5}\end{bmatrix}/\theta\begin{bmatrix}1\\ \frac{3}{5}\end{bmatrix}$ satisfies the following two algebraic equations:

$$f(x)=x^2\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(z)-x\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(z)-\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(z)=0,$$
$$g(x)=x^2\zeta_5^2\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(w)-x\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(w)+\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(w)=0.$$

Since $f(x)$ and $g(x)$ both have the same solution, it follows that

$$R(f,g)=\begin{vmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(z) & -\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(z) & -\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(z) & 0\\ 0 & \theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(z) & -\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(z) & -\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z)\\ \zeta_5^2\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(w) & -\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(w) & \theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(w) & 0\\ 0 & \zeta_5^2\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(w) & -\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(w) & \theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(w)\end{vmatrix}$$
$$=0,$$

which shows that

$$\left\{\zeta_5^3\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(w)+\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(w)\right\}^2$$
$$=\left\{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(z)\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(w)-\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(w)\right\}$$
$$\times\left\{\zeta_5^3\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(w)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(w)+\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(z)\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(w)\right\}.$$

Setting $z=w=0$, we have

$$\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\left\{\zeta_5^3\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}+\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}^2$$
$$=\left\{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}-\theta^3\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}\left\{\zeta_5^3\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}+\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}.$$

Let us show that

$$\zeta_5^3\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}+\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\not\equiv 0.$$

For this purpose, suppose that for every $\tau\in\mathbb{H}^2$,

$$\zeta_5^3\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)+\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)\equiv 0,$$

which implies that

$$\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)-\theta^3\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)\equiv 0,$$

or

$$\zeta_5^3\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}(0,\tau)+\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}(0,\tau)\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)\equiv 0.$$

Thus, it follows that

$$\zeta_5+\left(\frac{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}(0,\tau)}\right)^5\equiv 0,$$

which implies that $\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right](0,\tau)/\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right](0,\tau)$ is a constant. Taking the limit $\tau \longrightarrow i\infty$, we have

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{9}{5}\end{smallmatrix}\right](0,100\tau) = \exp\left(\frac{6\pi i}{50}\right)\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right](0,100\tau),$$

in which we used Jacobi's triple product identity (2.4). Since

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right](0,100\tau) = \sum_{n\in\mathbb{Z}}\exp(2\pi i)\left[\left(n+\frac{1}{10}\right)\frac{9}{10}\right]x^{(10n+1)^2}$$

and

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right](0,100\tau) = \sum_{n\in\mathbb{Z}}\exp(2\pi i)\left[\left(n+\frac{1}{10}\right)\frac{3}{10}\right]x^{(10n+1)^2},\ \ x=\exp(\pi i\tau),$$

comparing the coefficients of x^{121}, we have $\zeta_5^3 = 1$, which is impossible.

Therefore, we have

$$\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{1}{5}\end{smallmatrix}\right] = \frac{\left\{\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{7}{5}\end{smallmatrix}\right] - \theta^3\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{9}{5}\end{smallmatrix}\right]\right\}\left\{\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{7}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{9}{5}\end{smallmatrix}\right] + \zeta_5^3\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\right\}}{\left\{\zeta_5^3\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{7}{5}\end{smallmatrix}\right] + \theta\left[\begin{smallmatrix}\frac{1}{5}\\ 1\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{1}{5}\\ \frac{9}{5}\end{smallmatrix}\right]\right\}^2}.$$

□

6.3 The case for $j = 3$

Theorem 6.5. *One of* $\theta\left[\begin{smallmatrix}3/5\\ k/5\end{smallmatrix}\right], (k = 1, 3, 5, 7, 9)$, *is given by two rational expressions of the other four theta constants.*

For the proof, we obtain the following proposition and theorem, which can be proved in the same way as in the previous subsection.

Proposition 6.6. *For every* $(\zeta,\tau) \in \mathbb{C}\times\mathbb{H}^2$, *we have*

$$\begin{aligned}\theta^2\left[\begin{smallmatrix}1\\ \frac{3}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{3}{5}\\ \frac{1}{5}\end{smallmatrix}\right](\zeta,\tau)\theta\left[\begin{smallmatrix}\frac{3}{5}\\ \frac{3}{5}\end{smallmatrix}\right](\zeta,\tau) - \theta^2\left[\begin{smallmatrix}1\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}\frac{3}{5}\\ \frac{3}{5}\end{smallmatrix}\right](\zeta,\tau)\theta\left[\begin{smallmatrix}\frac{3}{5}\\ \frac{7}{5}\end{smallmatrix}\right](\zeta,\tau)\\ + \theta\left[\begin{smallmatrix}1\\ \frac{1}{5}\end{smallmatrix}\right]\theta\left[\begin{smallmatrix}1\\ \frac{3}{5}\end{smallmatrix}\right]\theta^2\left[\begin{smallmatrix}\frac{3}{5}\\ 1\end{smallmatrix}\right](\zeta,\tau) = 0,\end{aligned} \tag{6.16}$$

$$\zeta_5\theta^2\left[\begin{array}{c}1\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right](\zeta,\tau)-\zeta_5\theta^2\left[\begin{array}{c}1\\ \frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right](\zeta,\tau)$$
$$+\theta\left[\begin{array}{c}1\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}1\\ \frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right](\zeta,\tau)=0, \tag{6.17}$$

$$\theta^2\left[\begin{array}{c}1\\ \frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right](\zeta,\tau)+\zeta_5\theta^2\left[\begin{array}{c}1\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right](\zeta,\tau)$$
$$-\zeta_5\theta\left[\begin{array}{c}1\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}1\\ \frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right](\zeta,\tau)=0, \tag{6.18}$$

$$\theta^2\left[\begin{array}{c}1\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right](\zeta,\tau)+\zeta_5\theta^2\left[\begin{array}{c}1\\ \frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right](\zeta,\tau)$$
$$-\theta\left[\begin{array}{c}1\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}1\\ \frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right](\zeta,\tau)=0, \tag{6.19}$$

$$\theta^2\left[\begin{array}{c}1\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right](\zeta,\tau)-\theta^2\left[\begin{array}{c}1\\ \frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right](\zeta,\tau)\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right](\zeta,\tau)$$
$$+\zeta_5\theta\left[\begin{array}{c}1\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}1\\ \frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right](\zeta,\tau)=0. \tag{6.20}$$

Theorem 6.7. *For every $\tau\in\mathbb{H}^2$, we have*

$$\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]=\frac{\left\{\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]-\theta^3\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]\right\}\left\{\zeta_5^4\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]+\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]\right\}}{\left\{\zeta_5^4\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]+\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]\right\}^2}, \tag{6.21}$$

$$\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]=\frac{\left\{\zeta_5\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]+\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\right\}\left\{\theta^3\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]-\zeta_5^2\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]\right\}}{\left\{\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]+\zeta_5\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]\right\}^2}, \tag{6.22}$$

$$\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]=\frac{\left\{\zeta_5^4\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]+\theta^2\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]\right\}\left\{\theta^3\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]+\zeta_5^4\theta^3\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]\right\}}{\left\{-\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]+\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]\right\}^2}, \tag{6.23}$$

$$\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]=\frac{\left\{\zeta_5\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]+\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\right\}\left\{\theta\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]-\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]\right\}}{\left\{\theta^2\left[\begin{array}{c}\frac{3}{5}\\ 1\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{3}{5}\end{array}\right]+\zeta_5^4\theta^2\left[\begin{array}{c}\frac{3}{5}\\ \frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\ \frac{2}{5}\end{array}\right]\right\}^2}, \tag{6.24}$$

$$\theta\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]=\frac{\left\{\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]-\zeta_5^3\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{4}{5}\end{array}\right]\right\}\left\{-\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\zeta_5^2\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{4}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\right\}}{\left\{-\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]\right\}^2},\tag{6.25}$$

$$\theta\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]=\frac{\left\{-\zeta_5\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\zeta_5\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]\right\}\left\{-\zeta_5^2\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\right\}}{\left\{\zeta_5^2\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]-\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\right\}^2},\tag{6.26}$$

$$\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]=\frac{\left\{\zeta_5\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\right\}\left\{\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]+\zeta_5\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\right\}}{\left\{-\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]+\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\right\}^2},\tag{6.27}$$

$$\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{4}{5}\end{array}\right]=\frac{\left\{\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\zeta_5^4\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\right\}\left\{-\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\right\}}{\left\{\zeta_5\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\right\}^2},\tag{6.28}$$

$$\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]=\frac{\left\{\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]-\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\right\}\left\{\zeta_5\theta^2\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{4}{5}\end{array}\right]+\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\right\}}{\left\{\zeta_5\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]+\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\right\}^2},\tag{6.29}$$

$$\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]=\frac{\left\{\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]+\zeta_5^4\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\right\}\left\{-\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{1}{5}\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\zeta_5^2\theta\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\theta^3\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]\right\}}{\left\{\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]\theta^2\left[\begin{array}{c}\frac{3}{5}\\\frac{3}{5}\end{array}\right]+\zeta_5\theta^2\left[\begin{array}{c}\frac{3}{5}\\1\end{array}\right]\theta\left[\begin{array}{c}\frac{3}{5}\\\frac{2}{5}\end{array}\right]\right\}^2}.\tag{6.30}$$

7 Certain theta constant identities

In this section, for $j=1,3$, we investigate the algebraic structure of the following five theta constants:

$$\theta\left[\begin{array}{c}\frac{j}{5}\\\frac{k}{5}\end{array}\right],\ (k=1,3,5,7,9).$$

7.1 The case for $j=1$

In this subsection, we prove the following theorem:

Theorem 7.1. *Among the following five theta constants,*

$$\theta\begin{bmatrix}\frac{1}{5}\\ \frac{k}{5}\end{bmatrix},\ (k=1,3,5,7,9),$$

there exist at most three theta constants which are algebraically independent over $\mathbb{C}$.

For the proof, we obtain the following theorem:

Theorem 7.2. *For every* $\tau\in\mathbb{H}^2$, *we have*

$$\begin{aligned}
&\left\{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}+\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}^3\left\{\zeta_5\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}-\zeta_5\theta^3\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}\\
=&\left\{\zeta_5^3\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}+\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}^3\left\{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}-\zeta_5^4\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\},
\end{aligned}$$

$$\begin{aligned}
&\left\{\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}+\zeta_5^3\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\right\}^3\left\{\theta^3\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}+\zeta_5^3\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}\\
=&\left\{\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}-\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}^3\left\{\zeta_5^2\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}+\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\right\},
\end{aligned}$$

$$\begin{aligned}
&\left\{\zeta_5^4\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}-\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\right\}^3\left\{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}-\zeta_5\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\right\}\\
=&\left\{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}-\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\right\}^3\left\{\zeta_5\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}-\zeta_5^2\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\right\},
\end{aligned}$$

$$\begin{aligned}
&\left\{\zeta_5^2\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}+\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\right\}^3\left\{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}+\zeta_5^2\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}\\
=&\left\{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}-\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}\right\}^3\left\{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{9}{5}\end{bmatrix}+\zeta_5^3\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\right\},
\end{aligned}$$

$$\begin{aligned}
&\left\{\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}+\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\right\}^3\left\{\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}-\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\right\}\\
=&\left\{\zeta_5^2\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\right\}^3\left\{-\zeta_5^3\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}+\zeta_5^2\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta^3\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\right\}.
\end{aligned}$$

Proof. The first equation follows from equations (6.6) and (6.7). The second equation is obtained by equations (6.8) and (6.9). The third equation is derived from equations (6.10) and (6.11). The fourth equation follows from equations (6.12) and (6.13). The fifth equation follows from equations (6.14) and (6.15). □

7.2 The case for $j = 3$

In this subsection, we prove the following theorem:

Theorem 7.3. *Among the following five theta constants,*

$$\theta\begin{bmatrix} \frac{3}{5} \\ \frac{k}{5} \end{bmatrix},\ (k = 1, 3, 5, 7, 9),$$

there exist at most three theta constants which are algebraically independent over $\mathbb{C}$.

For the proof, we obtain the following theorem, which can be proved in the same way as in the previous subsection.

Theorem 7.4. *For every* $\tau \in \mathbb{H}^2$, *we have*

$$\begin{aligned}
&\left\{\theta\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\theta^2\begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} + \zeta_5\theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{2}{5} \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\right\}^3\left\{\zeta_5^3\theta\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{2}{5} \end{bmatrix} - \zeta_5^3\theta^3\begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\right\} \\
=&\left\{\zeta_5^4\theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{2}{5} \end{bmatrix} + \theta\begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}\theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\right\}^3\left\{\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - \zeta_5^2\theta\begin{bmatrix} \frac{3}{5} \\ \frac{2}{5} \end{bmatrix}\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\right\},
\end{aligned}$$

$$\begin{aligned}
&\left\{\theta^2\begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix} + \zeta_5^4\theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{1}{5} \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{2}{5} \end{bmatrix}\right\}^3\left\{\theta^3\begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{2}{5} \end{bmatrix} + \zeta_5^4\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{1}{5} \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\right\} \\
=&\left\{\theta\begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}\theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{2}{5} \end{bmatrix} - \theta\begin{bmatrix} \frac{3}{5} \\ \frac{1}{5} \end{bmatrix}\theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\right\}^3\left\{\zeta_5\theta\begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix} + \theta\begin{bmatrix} \frac{3}{5} \\ \frac{1}{5} \end{bmatrix}\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{2}{5} \end{bmatrix}\right\},
\end{aligned}$$

$$\begin{aligned}
&\left\{\zeta_5^2\theta\begin{bmatrix} \frac{3}{5} \\ \frac{7}{5} \end{bmatrix}\theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{9}{5} \end{bmatrix} - \theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{1}{5} \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\right\}^3\left\{\theta\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{9}{5} \end{bmatrix} - \zeta_5^3\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{1}{5} \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{7}{5} \end{bmatrix}\right\} \\
=&\left\{\theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{9}{5} \end{bmatrix} - \theta\begin{bmatrix} \frac{3}{5} \\ \frac{1}{5} \end{bmatrix}\theta^2\begin{bmatrix} \frac{3}{5} \\ \frac{7}{5} \end{bmatrix}\right\}^3\left\{\zeta_5^3\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{7}{5} \end{bmatrix}\theta\begin{bmatrix} \frac{3}{5} \\ \frac{9}{5} \end{bmatrix} - \zeta_5\theta\begin{bmatrix} \frac{3}{5} \\ \frac{1}{5} \end{bmatrix}\theta^3\begin{bmatrix} \frac{3}{5} \\ \frac{3}{5} \end{bmatrix}\right\},
\end{aligned}$$

$$\left\{\zeta_5\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}+\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\right\}^3\left\{\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}+\zeta_5\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\right\}$$
$$=\left\{\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}-\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\right\}^3\left\{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}+\zeta_5^4\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\right\},$$

$$\left\{\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}+\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\end{bmatrix}\right\}^3\left\{\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\end{bmatrix}-\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta^3\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\right\}$$
$$=\left\{\zeta_5\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\right\}^3\left\{-\zeta_5^4\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}+\zeta_5\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\theta^3\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\end{bmatrix}\right\}.$$

8 The algebraic structure of modular forms of level five

In this section, we investigate the algebraic structure of the following theta constants:

$$\theta\begin{bmatrix}\frac{j}{5}\\ \frac{k}{5}\end{bmatrix},\ \theta\begin{bmatrix}1\\ \frac{j}{5}\end{bmatrix},\ (j=1,3,k=1,3,5,7,9).$$

8.1 Certain theta constant identities

Theorem 8.1. *For every $\tau\in\mathbb{H}^2$, we have*

$$\left\{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}-\zeta_5^4\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}\right\}\left\{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}-\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\right\}$$
$$=\left\{\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}-\zeta_5^2\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\end{bmatrix}\right\}\left\{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}-\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\right\},\quad(8.1)$$

$$\left\{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\right\}\left\{\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}+\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\right\}$$
$$=\left\{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{7}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\right\}\left\{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}+\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\end{bmatrix}\right\},\quad(8.2)$$

$$\left\{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}-\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{9}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\right\}\left\{\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\\ \frac{7}{5}\end{bmatrix}+\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{9}{5}\end{bmatrix}\right\}$$
$$=\left\{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}-\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{9}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\right\}\left\{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\\ \frac{7}{5}\end{bmatrix}+\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{9}{5}\end{bmatrix}\right\}, \quad (8.3)$$

$$\left\{-\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{1}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\right\}\left\{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{1}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\right\}$$
$$=\left\{-\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{1}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{1}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\right\}\left\{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{3}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{1}{5}\end{bmatrix}\right\}, \quad (8.4)$$

$$\left\{\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{9}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\right\}\left\{\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\\ \frac{7}{5}\end{bmatrix}+\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{9}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\right\}$$
$$=\left\{\zeta_5\theta^2\begin{bmatrix}\frac{3}{5}\\ \frac{2}{5}\\ \frac{7}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{9}{5}\end{bmatrix}+\theta^2\begin{bmatrix}\frac{3}{5}\\ 1\end{bmatrix}\theta\begin{bmatrix}\frac{3}{5}\\ \frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\right\}\left\{\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{3}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ \frac{2}{5}\\ \frac{7}{5}\end{bmatrix}+\zeta_5^2\theta^2\begin{bmatrix}\frac{1}{5}\\ \frac{3}{5}\\ \frac{9}{5}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{5}\\ 1\end{bmatrix}\right\}. \quad (8.5)$$

Proof. Equation (8.1) follows from equations (4.6) and (4.16). Equation (8.2) is obtained by equations (4.7) and (4.17). Equation (8.3) is derived from equations (4.8) and (4.18). Equation (8.4) follows from equations (4.9) and (4.19). Equation (8.5) is obtained from equations (4.10) and (4.20). □

8.2 Proof of Theorem 8.2

Theorem 8.2. *Among the following twelve theta constants,*

$$\theta\begin{bmatrix}\frac{j}{5}\\ \frac{k}{5}\end{bmatrix},\ \theta\begin{bmatrix}1\\ \frac{j}{5}\end{bmatrix},\ (j=1,3,\ k=1,3,5,7,9),$$

there exist at most five theta constants which are algebraically independent over $\mathbb{C}$.

Proof. Theorem 8.2 follows from Theorems 5.1, 6.2, 6.5, 7.1, 7.3 and 8.1. □

References

[1] H. M. Farkas and I. Kra, *Automorphic forms for subgroups of the modular group.* Israel J. Math. **82**, (1993), 87-13.

[2] H. M. Farkas and I. Kra, *Theta Constants, Riemann Surfaces and the Modular Group.* AMS Grad Studies in Math, vol 37, 2001.

[3] H. M. Farkas, *Theta functions in complex analysis and number theory.* Surveys in number theory, Dev. Math., **17**, (2008), 57-87.

[4] K. Matsuda, *The determinant expressions of some theta constant identities.* Ramanujan J. **34**, (2014), 449-456.